

**WHAT IS CLAIMED IS:**

1. A method of determining prognosis of prostate cancer from a patient, comprising the steps of:

establishing the threshold value of at least one prognostic gene selected from the group of

5 Table 1A, 1B and 1C,

determining the amount of said prognostic gene from a prostate tissue of a patient,

comparing the amount of said prognostic gene with the established threshold value of said prognostic gene, and

determining the prognostic outcome of the patient.

10

2. The method according to Claim 1, wherein said prognostic gene is trp-p8.

3. The method according to Claim 1, wherein the amount of said prognostic gene is determined after amplification of said prognostic gene.

15

4. The method according to Claim 1, further comprising determining prostate-specific antigen.

5. The method of Claim 1, wherein said prognosis is determined after a selected therapeutic treatment.

20

6. The method according to Claim 1, wherein said prognosis contributes to selection of a therapeutic strategy.

7. The method according to Claim 1, wherein said prognosis contributes to selection of a therapeutic strategy.

25

8. A method of determining prognosis of prostate cancer from a patient, comprising the steps of:

establishing the threshold value of at least one prognostic gene selected from the group of  
Table 2,

determining the amount of said prognostic gene from a prostate tissue of a patient,  
comparing the amount of said prognostic gene with the established threshold value of

- 5 said prognostic gene, and  
determining the prognostic outcome of the patient.